Huntington's disease

S Davies, D B Ramsden

Abstract

The most recent findings in the elucidation of the molecular pathology of Huntington's disease are reviewed. Particular interest has been paid to the role of huntingtin and its associated proteins in excitotoxicity mediated via NMDA and kainate receptors.

(J Clin Pathol: Mol Pathol 2001;54:409–413)

Keywords: Huntington's disease; neurodegenerative disorder; Huntingtin; NMDA receptors; kainate receptors

This short review sets out to consider the most recent findings in the search to elucidate the pathological mechanisms underlying the fatal, neurodegenerative disorder known as Huntington's disease (HD), first described by George Huntington in 1872. HD involves the extrapyramidal motor system and is characterised by chorea, a dementia which is progressive, and other psychiatric symptoms. In terms of brain pathology, GABAergic medium spiny striatal neurones are most affected, resulting in atrophy of the caudate nucleus, putamen, and globus pallidus; there is also pronounced atrophy of the cerebral cortex.^{1,2} The neuronal loss in the striatum is accompanied by pronounced gliosis. The vulnerable striatal neurones contain enkephalin, dynorphin, and substance P, and primarily innervate the substantia nigra and globus pallidus. Medium sized spiny neurones containing somatostatin, neuropeptide Y, and NADPH diaphorase, in addition to cholinergic interneurons and parvalbumin containing GABAergic neurones, are relatively spared.³ In the cerebral cortex, large neurones in layer VI are the most affected, with smaller amounts of degeneration seen in layers III and V.^{2–4} Neurones may also be lost in the thalamus, zona reticulata of the substantia nigra, superior olive, lateral tuberal nucleus of the hypothalamus, and deep cerebellar nuclei.^{3–5} In exploring the sequence of neurodegeneration, Glass *et al* looked at receptor changes in the basal ganglia in the various stages of HD (summarised in

table 1). Their data indicate an overlapping pattern of neurodegeneration of GABAergic efferent projection neurones with the progress of the disease, accompanied by the loss of cannabinoid receptors throughout the basal ganglia.⁶

The importance of CAG repeats

One of the important early triumphs of modern molecular biology has been the demonstration that the underlying cause of HD is the expansion of a CAG repeat sequence in the first exon of a gene on chromosome 4p16.3, which encodes the protein huntingtin.^{6–8} CAG is the codon for glutamine (Q in the single letter code for amino acids). The normal range for the number of Qs in the polyglutamine tract (polyQ) is between six and 34, with disease being found when polyQ is greater than 40. Interestingly, within the HD disease genome, expansion of trinucleotide repeats appears not to be limited solely to the huntingtin gene. Keckarevic *et al* reported that allelic frequency distributions for two other genes containing such repeat sequences (SCA1 and FRDA) were shifted towards larger alleles compared with healthy controls, and suggested that a common mechanism caused expansion in all three genes.⁹ The expanded CAG repeat is transmitted predominantly through the male germ line in humans and transgenic mice. Using the transgenic mouse model to explore transmission, Kovtun *et al* found that the size of the CAG repeat was influenced by the sex of the offspring from identical fathers. Thus, male offspring had expanded repeats, whereas the opposite was seen in females. This finding led them to propose that X and Y chromosomal encoded factors influenced embryonic DNA repair and replication.¹⁰ Furthermore, within striatal cells, expansion biased changes were shown to increase with age, suggesting that non-replication based mechanisms may also contribute to CAG repeat instability.¹¹

The pathology of HD

The first substantive step towards understanding the disease process was the demonstration that intrastriatal injections of kainic acid produced lesions similar to those seen in HD; the model was further refined by the use of quinolinic acid injections. This led to the suggestion that glutamatergic overactivity (excitotoxicity) plays a central role in the pathogenesis of HD. Subsequently, the excitotoxicity hypothesis has proved to be the dominant concept influencing thinking on the pathology of HD. Nevertheless, other pathways—oxidative stress, impaired energy metabolism, apoptosis, and abnormal protein–protein interactions—which in turn may cause transcriptional dysregulation and altered gene expression,^{4–12} have also attracted

Department of Medicine, Division of Medical Science, University of Birmingham, Birmingham B15 2TH, UK

S Davies
D B Ramsden

Correspondence to:
Dr Ramsden
D.B.Ramsden@bham.ac.uk

Accepted for publication
26 September 2001

Table 1 Receptor changes in Huntington's disease⁶

Decreased		Increased	
Receptor	Regions	Receptor	Region
Early clinical phase			
Cannabinoid CB (1)	CN, P, GPE	GABA (A)	GPE
Dopamine D2	CN, P, GPE		
Adenosine A (2a)	CN, P, GPE		
Intermediate clinical phases			
Cannabinoid CB (1)	CN, P, SN		
Dopamine D1	CN, P, SN		
Final clinical phase			
Cannabinoid CB (1)	CN, P, GPI	GABA (A)	GPI
Dopamine D1	CN, P, GPI		

CN, caudate nucleus; GPE, globus pallidus externum; GPI, globus pallidus internum; P, putamen; SN, substantia nigra.

much interest. Central to all these have been attempts to understand the role of huntingtin in normal physiology and in HD.

The role of huntingtin in HD

Huntingtin has no great structural homology with other human proteins, so determination of its normal function has proved difficult. Changes seen in embryonic stem cells of huntingtin knockout mice indicate that it is essential for normal nuclear and perinuclear organelles and that one factor regulating its synthesis is iron.¹³ The protein (~ 349 kDa) is composed of approximately 3150 amino acids (depending on the size of the polyQ tract).¹⁴ As with many other large proteins, huntingtin is made up of repeated structures. The polyQ tract starts at residue 18. In addition, there are 10 Huntingtin elongation factor 3 (a subunit of protein phosphatase 2A-TOR1 (HEAT)) repeats, each of which is a loosely conserved repeating segment of approximately 40 amino acids, forming two hydrophobic α helices and a polymorphic polyproline region.¹⁵ The protein is widely distributed throughout the central nervous system (CNS), and this has posed the question of how its altered functioning gives rise to selective neuronal loss in HD. Initially, the most likely explanation was thought to be that the huntingtin gene mutations confer a “deleterious gain of function”, with the concept of loss of function gaining ground more recently,¹⁶ although how the variant huntingtins go on to cause neurodegeneration has remained a mystery.

HUNTINGTIN AND THE FORMATION OF NUCLEAR AGGREGATES

Both normal and variant huntingtins are localised chiefly in the cytoplasm of neurones.¹⁷ The wild-type protein is associated with cytoplasmic granules, organelles involved in retrograde transport and protein degradation, in some cortical and striatal neurones.¹⁸ In HD, ubiquitinated, polyQ containing, proteolytic N-terminal fragments of huntingtin form insoluble deposits both in the cytoplasm and nucleus. Both transgenic animals and genetically modified cell lines have been used to investigate the effects of expanded polyQ repeats on the localisation and processing of huntingtin. In knockin mice, increasing polyQ size was shown to increase nuclear localisation of full length huntingtin, in addition to the formation of N-terminal inclusions and aggregates in medium spiny neurones,¹⁹ and, in striatal embryonic cell lines, localisation in nuclear and perinuclear organelles.²⁰ The formation of N-terminal nuclear aggregates causes a shift in the core catalytic component of the proteosome to the nucleus. Thus, the formation of nuclear aggregates reduces the availability of the proteosome for digestion of other intracellular proteins that need to be removed, such as the apoptosis promoting protein p53. Nevertheless, despite the increased attraction of the proteosome to the aggregates, the rate of proteolysis of the N-terminal fragments was inversely related to polyQ length.²¹ Thus, disturbed proteolysis is an important feature of

HD. However, it is unclear whether the formation of aggregates per se is the essential cytotoxic step or a consequence of cellular dysfunction. Inhibition of aggregate formation by molecular chaperones that localise with the aggregates, such as heat shock protein 70, reduced cell death, suggesting a direct causal relation between aggregate formation and cell death,^{22–23} although other data indicate that the formation of nuclear inclusions and apoptosis are not closely related, with huntingtin acting via other routes.²⁴ Huntingtin has several protease susceptible domains. Within the human CNS, different patterns of proteolysis were seen in different regions. In the cortex, cleavage at two N-terminal sites (termed A and B) predominated, whereas in the striatum the dominant cleavages occurred at one N-terminal (A) and a C-terminal site.²⁵ Huntingtin is a caspase substrate. Inhibition of digestion of huntingtin by caspases 3 and 6 reduced apoptotic rates in cell models, suggesting that caspase inhibitors might provide a potential new therapeutic approach to the treatment of the disease.^{26–27}

HUNTINGTIN, EXCITOTOXICITY, AND APOPTOSIS

Although the above evidence supports the concept that aggregate formation is a causal agent in neuronal death, it is unclear whether this is the sole mechanism and how variant huntingtins relate to other features of the pathology, such as excitotoxicity. Much has been written about excitotoxicity via the NMDA and kainate receptors in HD (recent, extensive reviews on the structures and functions of these are by Chittajallu *et al*,²⁸ Lerma *et al*,²⁹ Cull-Candy *et al*,³⁰ and Ravenscroft and Brothie³¹). Over the past two years, much of the intracellular signalling cascade underlying excitotoxicity in HD has been pieced together. Key players in this are the NMDA and kainate receptors, a protein with the formidable name of “post-synaptic density protein 95” (also known as synapse associated protein 90), abbreviated here to PSDP-95, mixed lineage kinase (MLK), and c-Jun-N-terminal kinase (JNK). Glutamate receptor stimulation was shown previously to be involved in MLK-I activation,³² and in a separate study JNK activation was linked to apoptosis.³³ Now the connection between these processes and the roles of huntingtin and PSDP-95 in excitotoxicity are becoming more clearly understood. PSDP-95 is a scaffold protein, which binds to several intracellular proteins and possesses guanylate cyclase activity. PSDP-95 binds via repeat units in its structure, termed PDZ domains, to the NMDA receptor. PSDP-95 plays a pivotal role in regulating synaptic plasticity and synaptogenesis.³⁴ Specifically, PSDP-95 binds to the NR2 NMDA receptor and kainate receptor, GluR6, subunits. Variant huntingtins with expanded polyQ tracts interfere with the binding of PSD-95 to NMDA and kainate glutamate receptors, causing both receptors to become hypersensitive,^{35–36} thereby allowing increased Ca^{2+} influx via specific uptake pathways.³⁷ In turn, the MLK isoform, MLK-2,³⁸ is activated, causing activation of MAP kinase kinases 4 and

7 and stress signalling kinase, SEK-1, and consequently JNK-2³⁹ (dominant negative SEK-1 blocks JNK-2 activation³⁹). Activated JNK2 phosphorylates the N-terminal region of c-Jun, which is one half of the transcription factor AP-1, thus modulating gene transcription. Activated JNK also phosphorylates the C-terminal region of MLK-2. This step appears to be crucial in the triggering of apoptotic cell death. Cotransfection of a dominant negative MLK-2 blocks apoptotic cell death induced by variant huntingtins.⁴⁰

Interestingly, full length variant huntingtins, rather than N-terminal fragments, were found to be potent triggers of apoptosis in model cells expressing the NMDA receptor subtype prevalent in human striatal medium spiny neurones. Therefore, both full length variant huntingtins and nuclear N-terminal fragment aggregates cause cell death by different mechanisms. The actions of the full length variants may account for the observation that cellular dysfunction can be seen before aggregate formation in the nucleus.⁴¹

MULTIPLE PATHWAYS LEADING TO CELL DEATH

There is continuing evidence that variant huntingtins affect multiple pathways, which potentially lead to cell death. One such pathway is the deprivation of neurotrophic support. Brain derived neurotrophic factor (BDNF) expression is reduced in the caudate and putamen of patients with HD.⁴² That enhanced expression of neurotrophic factors may mitigate the effects of neurotoxins and thus be a potential therapeutic strategy was explored in animal and cell models. Both BDNF, nerve growth factor 3 (NGF-3), and NGF4/5 support proved effective treatments against quinolinic acid treatment in the various models, with BDNF being the most effective, opening the possibility of treatment by implantation of neurotrophic factor releasing systems.^{42 43}

One potential cause of cell death that has received much interest in several neurodegenerative diseases, including HD, is oxidative stress. However, a detailed investigation of oxidative stress indices in the caudate nucleus, putamen, and frontal cortex of HD brains revealed no alterations in concentrations of thiobarbituric acid-malondialdehyde adducts and lipid peroxides (measures of lipid peroxidation), or of oxidative DNA and protein damage. These data strongly suggest that oxidative stress is not a major contributor to neurodegeneration in HD.⁴⁴ However, evidence indicates that an alternative marker of oxidation—isoprostane concentration—is increased. This has been reviewed by Greco *et al.*⁴⁵

Reduced energy production as a consequence of defective mitochondrial complex II function is a well established feature of the late stages of HD. Garseth *et al* presented evidence of reduced lactate and citrate in the cerebrospinal fluid (CSF) of patients with HD, and this led them to suggest that the decreased citrate values were indicative of astrocytic and neuronal dysfunction.⁴⁶ Whether the energy deficit is a cause or consequence of the disease

is uncertain. Using a ³¹P nuclear magnetic resonance assay, Lodi *et al* found that the phosphocreatine to inorganic phosphate ratio was reduced in muscle tissue of both symptomatic and presymptomatic gene positive individuals, indicating that impaired energy production is an early feature of the disease.⁴⁷ In contrast, Guidetti *et al* found no evidence of change in the functioning of mitochondrial complexes I to IV in the neostriatum and cortex of presymptomatic individuals and patients with mild symptoms.⁴¹ Guidetti *et al* also found no change in the activities of complexes I to IV in the striatum and cortex of transgenic mice expressing full length polyQ expanded huntingtin before changes in the morphology of striatal and cortical neurones. They concluded that the changes in energy metabolism were a consequence rather than a cause of the disease.⁴¹ However, another group using a different animal model—3-nitropropionic acid treatments—came to conclusions suggesting that energy impairment was at the root of the disease. Calabresi *et al* found that treatment with 3-nitropropionic acid, which is a specific inhibitor of complex II, produced long term NMDA mediated excitation in striatal medium spiny neurones.⁴⁸ This effect was dopamine dependent, with dopamine acting via D2 receptors, whereas it was negatively regulated via D1 receptors, all of which mirrors the human disease.²¹ In a quinolinic acid treated rat model, energy metabolism, as measured by 2-deoxy-(¹⁸F)fluoro-D-glucose uptake was impaired within one week of lesioning, and this impairment deepened with time. However, D2 receptor expression did not simply follow the changes in glucose analogue uptake. One week after lesioning D2 receptor levels increased, after which there was a progressive decline.⁴⁹ These last two sets of findings raise the question of the relation of animal models to the human condition. They clearly demonstrate changes that parallel pathological changes seen in humans but leave the question of the ultimate cause of HD unanswered.

Another potential cause of cell dysfunction in HD is disruption of gene transcription by binding of variant huntingtins to transcription factors and intracellular signalling proteins. Of particular interest have been the interactions with the cAMP response element (CRE) binding protein and p53. N-terminal huntingtin fragments were shown to colocalise with both of these proteins in nuclear aggregates in cell culture and transgenic mouse models.^{50 51} This colocalisation within nuclear aggregates occurs because of interaction between the polyQ repeats in huntingtin and a similar short repeat in the CRE binding protein.⁵¹

Protein kinase C (PKC) is thought to be important in long term potentiation affecting brain plasticity and learning. In symptomatic transgenic mice expressing polyQ, expanded huntingtin steady state levels of PKC beta II mRNA were decreased in brain compared with control animals. This parallels the reduction of this mRNA in the caudate-putamen of human HD brain. It was postulated that the reduction

in PKC may play a part in the reduced ability to store information in HD.⁵²

The multiplicity of other pathways potentially affected in HD is illustrated by the number of proteins recognised to interact with huntingtin. These are known either as huntingtin associated proteins (HAPs), huntingtin interaction proteins (HIPs), and huntingtin–yeast partners (HYPs). Examples are HAP-1, HYP-J (α adaptin C), and HIP-3, which are associated with vesicular trafficking. HYP-A and HYP-I have nuclear functions, HIP-2 is a ubiquitin conjugating enzyme, and HYP-F is part of the 26S proteasome. New members of these categories are still being discovered and the physiological functions of recognised ones elucidated. Most recently, the functions of HIP-1 have been clarified. This molecule is encoded by a large gene of 32 exons, scanning 215 kb, which gives rise to two transcripts.⁵³ The translation product is involved in D1 receptor trafficking, linking huntingtin and the endocytic protein, clathrin.⁵⁴ In an immortalised striatal neuronal cell line, dopamine altered the subcellular localisation of huntingtin, resulting in an increase in membrane fractions containing HIP-1 and clathrin. Another recently described member of the HIP family (the gene located on 12q24),⁵⁵ HIP-1R was also shown to be involved in endocytosis. The C-terminal, talin-like domain of this protein binds to F-actin, thus linking the endocytic machinery with the actin cytoskeleton of cells.⁵⁶ HIP-1 is also a pro-apoptotic protein, the over-expression of which results in rapid caspase 3 dependent cell death via the intrinsic pathway of apoptosis.⁵⁷ The association of huntingtin with HIP-1 attenuated the pro-apoptotic actions of HIP-1. Interestingly, wild-type huntingtin binds more strongly with HIP-1 than do the polyQ expanded forms, which thus represents another potential neurodegenerative mechanism in HD.

The latest HAPs and HIPs to be identified are HAP40 and HIP-12. HAP40 is a 40 kDa protein encoded within intron 22 of the factor VIII gene. In the presence of normal huntingtin, HAP 40 is located in the cytosol, and in its absence it is actively transferred to the nucleus, although the consequences of this are not known.⁵⁸ HIP-12 was discovered while elucidating the gene structure of HIP-1. HIP-12 has no detectable pro-apoptotic properties, does not bind to huntingtin, but does bind to HIP-1.⁵⁹

A model of cell death in neurodegenerative disease

Finally, in this brief review, mention should be made of an advance in the understanding of the rate of cell death in neurodegenerative diseases. Clarke *et al* studied photoreceptor decay in model systems and, with reference to HD, the uptake of 18F-glucose into the caudate nucleus of patients with HD, this being regarded as a measure of surviving neurones.⁶⁰ The mathematical description that best fitted the data was that of a constant risk of cell death, with the qualification that an exponential decrease in the risk of cell death may also be a valid

model. The constant risk model means that cell death would be random and that the death of a neurone would be independent of the fate of other neurones, whereas the exponential decrease model proposes that the chance of cell death decreases as the number of surviving neurones decreases. They propose that the features of a neurodegenerative disease such as HD—genotype dependence, the random nature of cell death within a specific population, the constant risk or exponentially decreasing risk of cell death, and the late onset of the clinical signs and symptoms, which clearly means that most neurones survive for a considerable period of time—imply that the organism exists in a steady state. This condition, which they term the mutant steady state, is different from the normal healthy state, and is achieved by adaptive changes in metabolism to cope with the effects of the mutant gene product.

- 1 Reddy PH, Williams M, Tagle DA. Recent advances in understanding the pathogenesis of Huntington's disease. *Trends Neurosci* 1999;22:248–55.
- 2 Sharp AH, Ross CA. Neurobiology of Huntington's disease. *Neurobiological Diseases* 1996;3:3–15.
- 3 Martin JB. Molecular basis of the neurodegenerative disorders. *N Engl J Med* 1999;340:1970–80.
- 4 Ho LW, Carmichael J, Swartz J, *et al*. The molecular biology of Huntington's disease. *Psychol Med* 2001;31:3–14.
- 5 Tobin AJ, Signer ER. Huntington's disease: the challenge for cell biologists. *Trends Cell Biol* 2000;10:531–6.
- 6 Glass M, Dragunow M, Faull RL. The pattern of neurodegeneration in Huntington's disease: a comparative study of cannabinoid, dopamine, adenosine and GABA(A) receptor alterations in the human basal ganglia in Huntington's disease. *Neuroscience* 2000;97:505–19.
- 7 Weatherall DJ, Ledingham JGG, Warrell DA, eds. Movement disorders. In: *Oxford textbook of medicine*, 3rd ed. Oxford: Oxford University Press, 1996:4012–14.
- 8 Gusella JF, Wexler NS, Conneally PM, *et al*. A polymorphic DNA marker genetically linked to Huntington's disease. *Nature* 1983;306:234–8.
- 9 Keckarevic D, Culjkovic B, Savic D, *et al*. The status of SCA1, MJD/SCA3, FRDA, DRPLA and MD triplet containing genes in patients with Huntington disease and healthy controls. *J Neurogenet* 2000;14:257–63.
- 10 Kovtun IV, Therneau TM, McMurray CT. Gender of the embryo contributes to CAG instability in transgenic mice containing a Huntington's disease gene. *Hum Mol Genet* 2000;9:2767–75.
- 11 Kennedy L, Shelbourne PF. Dramatic mutation instability in HD mouse striatum: does polyglutamine load contribute to cell-specific vulnerability in Huntington's disease? *Hum Mol Genet* 2000;9:2539–44.
- 12 Cha JJ. Transcriptional dysregulation in Huntington's disease. *Trends Neurosci* 2000;23:387–92.
- 13 Hilditch-Maguire P, Trettel F, Passani LA, *et al*. Huntingtin: an iron-regulated protein essential for normal nuclear and perinuclear organelles. *Hum Mol Genet* 2000;9:2789–97.
- 14 The Huntington's Disease Collaborative Research Group. A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes. *Cell* 1993;72:971–83.
- 15 Andrade MA, Bork P. HEAT repeats in Huntington's disease protein. *Nat Genet* 1995;11:115–16.
- 16 Cattaneo E, Rigamonti D, Golferdo D, *et al*. Loss of normal Huntingtin function: new developments in Huntington's disease research. *Trends Neurosci* 2001;24:182–8.
- 17 DiFiglia M, Sapp E, Chase K, *et al*. Huntingtin is a cytoplasmic protein associated with vesicles in human and rat brain neurons. *Neuron* 1995;14:1075–81.
- 18 Petersen A, Mani K, Brudin P. Recent advances on the pathogenesis of Huntington's disease. *Exp Neurol* 1999;157:1–18.
- 19 Wheeler VC, White JK, Gutekunst CA, *et al*. Long glutamine tracts cause nuclear localization of a novel form of huntingtin in medium spiny striatal neurons in HdhQ92 and HdhQ111 knock-in mice. *Hum Mol Genet* 2000;9:503–13.
- 20 Trettel F, Rigamonti D, Hilditch-Maguire P, *et al*. Dominant phenotypes produced by the HD mutation in STHdh(Q111) striatal cells. *Hum Mol Genet* 2000;9:2799–809.
- 21 Jana NR, Zemskov EA, Wang Gh, *et al*. Altered proteasomal function due to the expression of polyglutamine-expanded truncated N-terminal huntingtin induces apoptosis by caspase activation through mitochondrial cytochrome c release. *Hum Mol Genet* 2001;10:1049–59.
- 22 Jana NR, Tanaka M, Wang Gh, *et al*. Polyglutamine length-dependent interaction of Hsp40 and Hsp70 family chaperones with truncated N-terminal huntingtin: their role in suppression of aggregation and cellular toxicity. *Hum Mol Genet* 2000;9:2009–18.

- 23 Carmichael J, Chatellier J, Woolfson A, *et al*. Bacterial and yeast chaperones reduce both aggregate formation and cell death in mammalian cell models of Huntington's disease. *Proc Natl Acad Sci U S A* 2000;**97**:9701–5.
- 24 Saudou F, Finkbeiner S, Devys D, *et al*. Huntingtin acts in the nucleus to induce apoptosis but death does not correlate with the formation of intranuclear inclusions. *Cell* 1998;**95**:55–66.
- 25 Mende-Mueller LM, Toneff T, Hwang SR, *et al*. Tissue-specific proteolysis of Huntingtin (htt) in human brain: evidence of enhanced levels of N- and C-terminal htt fragments in Huntington's disease striatum. *J Neurosci* 2001;**21**:1830–7.
- 26 Wellington CL, Leavitt BR, Hayden MR. Huntington disease: new insights on the role of huntingtin cleavage. *J Neural Transm Suppl* 2000;**58**:1–17.
- 27 Ona V, Li M, Vonsattel J, *et al*. Inhibition of caspase-1 slows disease progression in a mouse model of Huntington's disease. *Nature* 1999;**399**:204–5.
- 28 Chittajallu R, Braithwaite SP, Clarke VR, *et al*. Kainate receptors: subunits, synaptic localization and function. *Trends Pharmacol Sci* 1999;**20**:26–35.
- 29 Lerma J, Paternain AV, Rodriguez-Moreno A, *et al*. Molecular physiology of kainate receptors. *Physiol Rev* 2001;**81**: 971–98.
- 30 Cull-Candy S, Brickley S, Farrant M. NMDA receptor subunits: diversity, development and disease. *Curr Opin Neurobiol* 2001;**11**:327–35.
- 31 Ravenscroft P, Brothie J. NMDA receptors in the basal ganglia. *J Anat* 2000;**196**:577–85.
- 32 Schwarzschild MA, Cole RL, Hyman SE. Glutamate, but not dopamine, stimulates stress-activated protein kinase and AP-1-mediated transcription in striatal neurons. *J Neurosci* 1997;**17**:3455–66.
- 33 Yang DD, Kuan CY, Whitmarsh AJ, *et al*. Absence of excitotoxicity-induced apoptosis in the hippocampus of mice lacking the Jnk3 gene. *Nature* 1997;**389**:865–70.
- 34 Che YH, Tamatani M, Tohyama M. Changes in mRNA for post-synaptic density-95 (PSD-95) and carboxy-terminal PDZ ligand of neuronal nitric oxide synthase following facial nerve transection. *Brain Res Mol Brain Res* 2000;**76**: 325–35.
- 35 Sun Y, Savanenin A, Reddy PH, *et al*. Polyglutamine-expanded huntingtin promotes sensitization of N-methyl-D-aspartate receptors via post-synaptic density 95. *J Biol Chem* 2001;**276**:24713–18.
- 36 Savinainen A, Garcia EP, Dorow D, *et al*. Kainate receptor activation induces mixed lineage kinase-mediated cellular signaling cascades via post-synaptic density protein 95. *J Biol Chem* 2001;**276**:11382–6.
- 37 Sattler R, Charlton MP, Hafner M, *et al*. Distinct influx pathways, not calcium load, determine neuronal vulnerability to calcium neurotoxicity. *J Neurochem* 1998;**71**: 2349–64.
- 38 Liu YF, Dorow D, Marshall J. Activation of MLK2-mediated signaling cascades by polyglutamine-expanded huntingtin. *J Biol Chem* 2000;**275**:19035–40.
- 39 Liu YF. Expression of polyglutamine-expanded Huntingtin activates the SEK1–JNK pathway and induces apoptosis in a hippocampal neuronal cell line. *J Biol Chem* 1998;**273**: 28873–7.
- 40 Phelan DR, Price G, Liu YF, *et al*. Activated JNK phosphorylates the c-terminal domain of MLK2 that is required for MLK2-induced apoptosis. *J Biol Chem* 2001;**276**:10801–10.
- 41 Guidetti P, Charles V, Chen EY, *et al*. Early degenerative changes in transgenic mice expressing mutant huntingtin involve dendritic abnormalities but no impairment of mitochondrial energy production. *Exp Neurol* 2001;**169**:340–50.
- 42 Ferrer I, Goutan E, Marin C, *et al*. Brain-derived neurotrophic factor in Huntington disease. *Brain Res* 2000;**866**:257–61.
- 43 Menei P, Pean JM, Nerriere-Daguin V, *et al*. Intracerebral implantation of NGF-releasing biodegradable microspheres protects striatum against excitotoxic damage. *Exp Neurol* 2000;**161**:259–72.
- 44 Alam ZI, Halliwell B, Jenner P. No evidence for increased oxidative damage to lipids, proteins, or DNA in Huntington's disease. *J Neurochem* 2000;**75**:840–6.
- 45 Greco A, Minghetti L, Levi G. Isoprostanes, novel markers of oxidative injury, help understanding the pathogenesis of neurodegenerative diseases. *Neurochem Res* 2000;**25**:1357–64.
- 46 Garseth M, Sonnewald U, White LR, *et al*. Proton magnetic resonance spectroscopy of cerebrospinal fluid in neurodegenerative disease: indication of glial energy impairment in Huntington chorea, but not Parkinson disease. *J Neurosci Res* 2000;**60**:779–82.
- 47 Lodi R, Schapira AH, Manners D, *et al*. Abnormal in vivo skeletal muscle energy metabolism in Huntington's disease and dentatorubropallidoluysian atrophy. *Ann Neurol* 2000;**48**:72–6.
- 48 Calabresi P, Gubellini P, Picconi B, *et al*. Inhibition of mitochondrial complex II induces a long-term potentiation of NMDA-mediated synaptic excitation in the striatum requiring endogenous dopamine. *J Neurosci* 2001;**21**: 5110–20.
- 49 Araujo DM, Cherry SR, Tatsukawa KJ, *et al*. Deficits in striatal dopamine D(2) receptors and energy metabolism detected by in vivo microPET imaging in a rat model of Huntington's disease. *Exp Neurol* 2000;**166**:287–97.
- 50 Steffan JS, Kazantsev A, Spasic-Boskovic O, *et al*. The Huntington's disease protein interacts with p53 and CREB-binding protein and represses transcription. *Proc Natl Acad Sci U S A* 2000;**97**:6763–8.
- 51 Nucifora FC, Jr, Sasaki M, Peters MF, *et al*. Interference by huntingtin and atrophin-1 with cbp-mediated transcription leading to cellular toxicity. *Science* 2001;**291**:2423–8.
- 52 Harris AS, Denovan-Wright EM, Hamilton LC, *et al*. Protein kinase C beta II mRNA levels decrease in the striatum and cortex of transgenic Huntington's disease mice. *J Psychiatry Neurosci* 2001;**26**:117–22.
- 53 Fichter KM, McCutcheon K, Drubin D, *et al*. HIP12 is a non-proapoptotic member of a gene family including HIP1, an interacting protein with huntingtin. *Mamm Genome* 2000;**11**:1006–15.
- 54 Kim M, Velier J, Chase K, *et al*. Forskolin and dopamine D1 receptor activation increase huntingtin's association with endosomes in immortalized neuronal cells of striatal origin. *Neuroscience* 1999;**89**:1159–67.
- 55 Seki N, Muramatsu M, Sugano S, *et al*. Cloning, expression analysis, and chromosomal localization of HIP1R, an isolog of huntingtin interacting protein (HIP1). *J Hum Genet* 1998;**43**:268–71.
- 56 Engqvist-Goldstein AE, Kessels MM, Chopra VS, *et al*. An actin-binding protein of the Sla2/Huntingtin interacting protein 1 family is a novel component of clathrin-coated pits and vesicles. *J Cell Biol* 1999;**147**:1503–18.
- 57 Hackam AS, Yassa AS, Singaraja R, *et al*. Huntingtin interacting protein 1 induces apoptosis via a novel caspase-dependent death effector domain. *J Biol Chem* 2000;**275**: 41299–308.
- 58 Peters MF, Ross CA. Isolation of a 40-kDa Huntingtin-associated protein. *J Biol Chem* 2001;**276**:3188–94.
- 59 Chopra VS, Metzler M, Rasper DM, *et al*. HIP12 is a non-proapoptotic member of a gene family including HIP1, an interacting protein with huntingtin. *Mamm Genome* 2000;**11**:1006–15.
- 60 Clarke G, Collins RA, Leavitt BR, *et al*. A one-hit model of cell death in inherited neuronal degenerations. *Nature* 2000;**406**:195–9.