

Alterations in p53 predict response to preoperative high dose chemotherapy in patients with gastric cancer

F Bataille, P Rümmele, W Dietmaier, D Gaag, F Klebl, A Reichle, P Wild, F Hofstädter, A Hartmann

J Clin Pathol: Mol Pathol 2003;**56**:286–292

Aims: To evaluate the usefulness of molecular markers in predicting histopathological and clinical response to preoperative high dose chemotherapy (HDCT) and survival of patients with advanced gastric cancer.

Methods: In a phase II trial, 25 patients with metastatic gastric cancer received preoperative tandem HDCT consisting of etoposide, cisplatin, and mitomycin, followed by autologous bone marrow transplantation to achieve surgical resectability. Samples before and after treatment, from normal and tumour tissue, were characterised histopathologically, and both p53 and BAX expression was analysed by immunohistochemistry. Pretreatment formalin fixed, paraffin wax embedded samples from normal and tumour tissue were microdissected, and the extracted DNA was preamplified using improved primer extension preamplification polymerase chain reaction. Detection of microsatellite instability (MSI) or loss of heterozygosity (LOH) was performed using markers for p53, BAX, BAT25, BAT26, D2S123, D17S250, and APC. Exons 5–9 of the p53 gene were sequenced directly on ABI 373.

Results: Four parameters were significantly associated with response to chemotherapy and prolonged overall survival: positive p53 immunostaining, positive p53 mutation status before chemotherapy, strong histological regression induced by preoperative HDCT, and surgical treatment. Patients' sex or age, tumour location or stage, lymph node status, Lauren classification, MSI, or LOH did not influence duration of survival significantly in this high risk population.

Conclusion: Positive p53 immunostaining and p53 mutation status in pretreatment tumour biopsies might be useful molecular predictors of response and prognosis in patients with advanced gastric cancer treated by preoperative HDCT.

See end of article for authors' affiliations

Correspondence to:
Dr F Bataille, Department of Pathology, University of Regensburg, 93042 Regensburg, Germany; frau.k.bataille@klinik.uni-regensburg.de

Accepted for publication 25 June 2003

Despite its decreasing prevalence, gastric carcinoma is still one of the major causes of cancer death worldwide.¹ Because most tumours are diagnosed late in locally advanced stages, the overall median survival is only 16 months.² Surgical treatment results in poor survival rates in advanced stages. Several studies have shown that preoperative chemotherapy can achieve a response in 50–60% of patients, and allows radical surgery in 40–50% of cases with previously unresectable tumours.^{3–8} To increase response rates and survival time, the benefit of high dose chemotherapy (HDCT) protocols with autologous bone marrow transplantation (ABMT) were evaluated for several cancer entities. HDCT is based on the hypothesis that increased dosage will overcome drug resistance, eradicate metastatic disease, and increase cure rates. Because of the notable toxicity of this treatment and the limited survival time of patients with advanced gastric cancer, it would be useful to be able to select those patients whose tumours will be sensitive to chemotherapy.^{9–11} Therefore, predictors for response to HDCT and survival would be clinically useful.

“Several studies have shown that preoperative chemotherapy can achieve a response in 50–60% of patients, and allows radical surgery in 40–50% of cases with previously unresectable tumours”

Clinical or histopathological parameters may be suitable. In addition, the characterisation of molecular alterations in tumour tissue may be useful to help predict response to chemotherapy.

The tumour suppressor protein p53 is a potent activator of apoptosis, and p53 dependent apoptosis modulates the

cytotoxic effects of common antitumour agents such as 5-fluorouracil, doxorubicin, and cisplatin.¹² p53 independent¹³ regulation of cisplatin induced tumour cell death includes expression of proteins of the Bcl-2 family,¹⁴ or activation of stress kinase cascades^{15–16} at the level of Bcl-2. Many apoptotic pathways converge, and the ratio of Bcl-2 to BAX protein might be the final determinant of whether a cell enters the execution phase.¹⁷ BAX is a dominant inhibitor of Bcl-2.¹⁸

It is well established that mismatch repair proteins are involved in triggering an apoptotic response after cisplatin damage.¹⁹ The mismatch repair protein complex can recognise and bind to 1,2–GpG adducts.^{20–22} Cells defective for mismatch repair become more resistant to G2 arrest and apoptosis provoked by cisplatin.^{23–26}

The aim of our study was to assess the clinical, histopathological, and molecular parameters of patients with advanced gastric cancer treated by HDCT/ABMT, and their association with response to treatment and survival.

MATERIAL AND METHODS

Patients

Between October 1996 and December 2000, 25 patients with locally advanced gastric carcinoma were treated with a preoperative HDCT protocol followed by ABMT in a phase II study. All patients were deemed unresectable before the start of treatment, and tissue blocks from all patients were available

Abbreviations: ABMT, autologous bone marrow transplantation; HDCT, high dose chemotherapy; LOH, loss of heterozygosity; MSI, microsatellite instability; MSI-H, microsatellite unstable; PCR, polymerase chain reaction

for study. There were 11 women and 14 men, with a mean age of 50 years (range, 35–65).

A modified EAP schedule was used for induction treatment (doxorubicin, 40 mg/m²; etoposide, 120 mg/m²; cisplatin, 40 mg/m²), followed by granulocyte colony stimulating factor (filgrastim, Neupogen[®]) started at day 4 at a dose of 10 µ/kg body weight administered subcutaneously. Autologous peripheral blood stem cells were collected by leukapheresis after leucocyte nadir (Cobe Spectra; Cobe Laboratories, Lakewood, Colorado, USA) on consecutive days. If patients had stable disease following induction treatment, the first cycle of HDCT (etoposide, 500 mg/m²; cisplatin, 50 mg/m²; mitomycin C, 10 mg/m²; and BCNU (nitroso urea), 300 mg/m²) was administered three weeks thereafter. Patients who achieved a reduction in tumour size of > 50% after the first cycle of HDCT qualified for an identical second cycle four weeks later.

All patients were clinically judged as cT3 or cT4 according to the TNM classification,²⁷ and had systemic metastasis. None of the patients had received systemic chemotherapy for gastric cancer previously. Response to chemotherapy was assessed by endoscopy, endosonography, and computed tomography scan, as defined by the Eastern Cooperation Oncology Group.²⁸

Resectability and clinical response were defined as present if patients achieved partial remission (> 50% size reduction of the target lesion).

Definitive clinical TNM staging yielded 19 patient with T4 tumours and six with T3 tumours. N1, N2, and N3 disease was found in nine, three, and 13 cases, respectively. No patient showed N0 disease.

Response to chemotherapy enabled the resection of the tumour in 16 patients. Local (R0) resection was possible in three patients with T3, and nine with T4. In three patients with T4 tumours, resection margins were infiltrated by the tumour (R1 resection), and one patient with T3 disease underwent palliative resection only. All nine non-responders underwent non-curative palliative procedures.

Survival time was defined as duration between the date of first chemotherapy and the date of death or last follow up (table 1). Median survival of the studied 25 patients was 13 months. Median survival for patients with response to treatment and subsequent gastrectomy was 16 months (range, three to 50) and 7.5 months (range, five to 11.5) for those without resection. Twelve patients from the first group survived more than 12 months, and four are still alive.

Table 1 summarises the clinical and histopathological data of all the patients.

Histopathological evaluation

Pretreatment biopsies and post-treatment specimens (16 gastrectomy specimens and nine diagnostic biopsies from non-resected patients after last chemotherapy) were evaluated by two pathologists (FB and PR). All tissue was fixed in 4% buffered formalin and embedded in paraffin wax. Histological sections were stained with haematoxylin and eosin; tumours were graded according to the World Health Organisation system³¹ and histologically classified according to the Lauren classification (table 1).²⁹

The gastrectomy specimens or post-treatment biopsies (five biopsies with at least 18 sections from different tumour areas from each non-resected patient) were evaluated for histological changes resulting from preoperative chemotherapy according to the Japanese Research Society for Gastric Cancer classification.³⁰

Immunohistochemistry

Immunohistochemical studies for the expression of p53 and BAX were performed on pretreatment tissue samples using an avidin–biotin peroxidase method with diaminobenzidine as chromogen. After antigen retrieval (microwave treatment of formalin fixed, paraffin wax embedded, 5 µm thick tissue sections for 40 minutes at 240 W in citrate buffer, pH 6.0), immu-

nohistochemistry was carried out in a NEXES immunostainer (Ventana Medical System, Tucson, Arizona, USA), according to the manufacturer's instructions. For p53, a mouse monoclonal antibody (Bp53–12; Santa Cruz Biotechnology, Santa Cruz, California, USA) was used at a dilution of 1/1000 as the primary antibody and for BAX immunostaining, a rabbit polyclonal IgG primary antibody (clone P-19; rabbit polyclonal IgG; Santa Cruz Biotechnology) was used. A positive reaction for p53 protein was defined as nuclear staining in > 10% of the tumour cell nuclei. A tumour was regarded as expressing BAX when more than > 5% of tumour cells had cytoplasmic staining.

The staining results were interpreted by one pathologist (FB) who was unaware of the clinical outcome of the patients.

Microdissection

Genomic DNA was prepared from 5 µm thick paraffin wax embedded sections of pretreatment tumour tissue after dewaxing and microdissection. Pure tumour cell populations were obtained using either manual microdissection with a needle under an inverted microscope (×40) or laser microdissection (PALM[®] Robot Microbeam; PALM, Wolfratshausen, Germany), as described previously.³²

DNA isolation

For genetic analysis, paired pretreatment normal and tumour DNA samples were extracted using the QIAmp DNA mini kit (Qiagen, Hilden, Germany), according to the protocol provided by the manufacturer. The purified DNA template was amplified by whole genome amplification³³ before further polymerase chain reaction (PCR) analyses.³⁴

Microsatellite analysis

Microsatellite analysis was performed using the recommended reference panel for detection of MSI in colorectal cancer. This panel is composed of two mononucleotide repeats (BAT25 and BAT26) and three dinucleotide repeats (D5S346, D2S123, and D17S250).³⁵ PCR amplification was performed as described previously.³⁶ Subsequently, PCR products were analysed by 6.7% polyacrylamide/50% urea gel electrophoresis in a SequiGen sequencing gel chamber (BioRad, Hercules, California, USA), followed by silver nitrate staining.³⁷

All gels were evaluated independently by two observers (WD and FB). A tumour was classified as microsatellite unstable (MSI-H) if two or more markers showed instability.

LOH analysis (p53 ALK, D2S123, D17S250, and APC) was performed as described previously.³⁸ Informative cases were scored as allelic losses when the intensity of a signal for an allele from tumour DNA was decreased to 50%, relative to the matched allele from normal DNA. All cases of LOH and MSI were verified in an independent PCR reaction.

p53 mutation

Exons 5–9 of the p53 tumour suppressor gene were sequenced directly using single exon amplification using an ABI 373 sequencer, as described previously.³⁴

Statistics

Survival curves were constructed using the Kaplan–Meier method and differences were assessed by the log rank test. The χ^2 test or Fisher's exact test (two sided) was performed to determine the correlation between molecular features, clinicopathological parameters, and the survival of patients. A p value of < 0.05 was considered to be significant. All statistical analyses were conducted using SPSS software (SPSS, Chicago, Illinois, USA).

RESULTS

Parameters associated with response to chemotherapy and survival were positive p53 immunohistochemistry

Table 1 Clinical and histopathological data, results of immunohistochemistry, p53 sequencing, and microsatellite analyses in 25 advanced gastric carcinomas

Clinical data																										
Patient ID	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	
Sex	f	f	m	m	f	m	f	f	m	f	m	f	m	f	m	m	f	f	f	m	m	m	m	m	m	m
Age at diagnosis (years)	57	49	62	46	61	42	51	46	32	45	55	47	48	49	44	43	44	33	54	50	40	56	60	34	36	
Overall survival (months)	12.3	5.5	50.1	17.7	20.8	6.2	8.1	7.4	9.5	11.8	12.6	4.9	14.5	12.3	4.2	13.3	5	29.6	6.9	12.6	3	16.7	11.6	19.7	4.7	
Survival status	1	1	0	1	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	0	1	0	1	
Resection	1	0	1	1	1	0	0	0	0	1	1	1	1	1	1	1	0	1	0	1	1	1	0	1	0	
Histopathological data																										
Localisation	co	an	co/an	ca	co	co	an	an	an	co	an	co/an	co	co/an	an	ca	an	an	co	an	co	an	co	co	ca	
Lauren classification	d	d	d	i	i	d	d	i	d	d	d	d	i	i	d	i	d	d	d	d	d	i	d	d	d	
Histological regression																										
Regression score	0	0	3	3	3	0	0	0	0	3	2	1a	1b	2	1a	1b	0	3	0	2	2	3	0	3	0	
Immunohistochemistry																										
p53	1	0	1	1	1	0	0	1	0	1	1	1	1	0	0	0	0	1	0	1	0	1	1	1	0	
BAX	0	0	0	1	0	0	1	1	0	1	1	0	1	1	0	0	0	1	0	0	0	0	1	0	1	
Microsatellite analysis																										
<i>Mononucleotide markers</i>																										
BAT25																										
BAT26																										
<i>Dinucleotide markers</i>																										
APC				LOH	LOH		LOH					LOH	LOH			LOH			LOH	LOH	LOH					
D17S250																							LOH			
D2S123		MSI		MSI															MSI	MSI						
<i>Pentanucleotide marker</i>																										
TP53.Alk					LOH							LOH				LOH	LOH									
Sequencing results																										
p53 mutation	0	0	1	1	1	0	0	1	0	0	0	1	1	0	0	0	0	0	0	1	0	0	0	1	0	
Exon			5	8	5			8				6	5						6				5			
Codon			177	282	162			273				213	175						220				178			
Wild-type sequence			CCC	CGG	CGC			CGT				CGA	CGC						TAT				GCC			
Nucleotide substitution			TCC	CTG	CCC			TGT				CCA	CAC						TGT				GAC			
Allelic status of p53 mutation			het	het	het			het				het	het						het				het			

Sex: f, female; m, male. Survival status: 0, censored; 1, death. Localisation: ca, cardia; co, corpus; an, antrum. Histopathological classification according to Lauren²⁹: i, intestinal type; d, diffuse type. Regression score according to the Japanese Research Society for Gastric Cancer classification of gastric carcinoma³⁰: 0, grade 0; 1a, grade 1a; 1b, grade 1b; 2, grade 2; 3, grade 3. p53 immunohistochemistry: 0, negative; 1, positive (nuclear expression in >10% of tumour cells). BAX immunohistochemistry: 0, negative; 1, positive (cytoplasmatic expression in >5% of tumour cells). Allelic status of p53 mutation: het, heterozygeous; hemi, hemizygeous.

Table 2 Clinicopathological characteristics and their association with overall survival in 25 patients with metastatic gastric cancer

Variable	Coding plan	Overall survival				Log rank
		N	Events	Censored	Survival (months)	
Surgery	0 No surgery	9	8	1	7.5	0.0007
	1 Surgery	16	13	3	16.0	
Histological regression	0 No regression	14	13	1	8.2	0.0018
	1 Regression	11	8	3	18.8	
p53 mutation	0 No p53 mutation	17	15	2	10.2	0.0439
	1 p53 mutation	8	6	2	18.5	
p53 immunohistochemistry	0 Negative	11	10	1	7.2	0.0003
	1 Positive	14	11	3	17.3	

Regression score according to the Japanese Research Society for Gastric Cancer classification of gastric carcinoma³⁰: 0, grade 0; 1a, grade 1a; 1b, grade 1b; 2, grade 2; 3, grade 3. p53 immunohistochemistry: 0, negative; 1, positive (nuclear expression in >10% of tumour cells).

($p = 0.0003$), positive p53 mutation status ($p = 0.044$), moderate to severe histological regression after chemotherapy ($p = 0.002$), and surgical treatment ($p = 0.0007$) (table 2). No associations were seen between response or survival time and sex, age, tumour location and stage, lymph node status, or Lauren classification.

The histological effects of preoperative chemotherapy included degeneration of nuclei, coagulation necrosis of cancer cells, with or without formation of cholesterol crystals, appearance of giant cells with multiple nuclei, foamy degeneration of cancer cells, or creation of extracellular mucin pools without associated neoplastic cells (fig 1A–D).

Histologically, response (grade 2 and 3) could be verified in 11 of the 25 tumours. There was no association with the Lauren classification or degree of differentiation. Nine of 12

responders revealed histological regression ($p = 0.002$). Despite our small sample size in this study, pretreatment p53 alterations were significantly associated with response to HDCT and longer survival. A high degree of p53 staining was detected in pretreatment biopsies in 14 of 25 patients (fig 2B). There was a significant association between positive p53 immunostaining in pretreatment biopsies and a survival time longer than 12 months ($p = 0.0003$), pretreatment presence of p53 mutation ($p = 0.0001$), and histological regression ($p = 0.002$). p53 mutations were found in eight of the tumours (fig 2C,D), and all of them were associated with p53 overexpression ($p = 0.0001$). In six of them, survival time was longer than 12 months ($p = 0.044$).

Pretreatment positive BAX immunoreactivity was seen in 10 of the 25 patients (table 1; fig 2A). There was no significant




Figure 1 Histological characteristics of tumour regression after chemotherapy. (A) foamy degeneration of cancer cells (arrows); original magnification, $\times 200$; haematoxylin and eosin stained. (B) Coagulation necrosis of cancer cells with formation of cholesterol crystals (arrows); original magnification, $\times 100$; haematoxylin and eosin stained. (C) Neoplastic cells with stromal fibrosis; original magnification, $\times 200$; haematoxylin and eosin stained. (D) Residual neoplastic cells with stromal fibrosis (arrows); original magnification, $\times 400$; haematoxylin and eosin stained.




Figure 2 Immunohistochemical and molecular characterisation of pretreatment biopsy specimens in patients with gastric cancer. (A) Strong cytoplasmic BAX expression in most of the cancer cells; original magnification, $\times 630$. (B) Strong nuclear p53 expression in most of the cancer cells; original magnification, $\times 400$. (C) p53 mutation in patient 13, exon 5, codon 175, CGC \rightarrow CAC ($5' \rightarrow 3'$). (D) p53 mutation in patient 13, exon 5, codon 175, CGC \rightarrow CAC ($3' \rightarrow 5'$).

association with clinical and histological response to HDCT ($p = 0.215$).

Microsatellite analyses revealed low MSI (only one marker showing instability) in pretreatment biopsies in four of 25 tumours (table 1). None of the cases showed MSI-H or alterations at the mononucleotide markers BAT25 and BAT26.

LOH was detected most frequently at the APC locus (in nine of 25 patients), and the p53 locus (in four of 25 patient; table 1). Neither MSI nor LOH was associated with response to chemotherapy or survival.

DISCUSSION

Most patients with gastric cancer are unresectable at diagnosis, or will suffer a relapse after surgery, resulting in a five year overall survival of less than 20%.^{39–41} Preoperative chemotherapy could be an approach that might improve surgical resectability, which is one of the main prognostic factors in patients with gastric carcinoma.⁴² However, only 40–50% of patients benefit from this treatment modality, whereas approximately 30% of patients experience moderate to severe toxic side effects.^{6, 8, 43} Understanding the molecular genetic features that determine response or resistance to chemotherapy could permit the selection of the most suitable patients for preoperative treatment.

“Histological responders underwent surgical treatment in all cases and survived significantly longer than non-responders”

To test the hypothesis that dose escalation would provide a benefit with regard to resectability and overall survival, a phase II study was designed to determine the feasibility of etoposide, cisplatin, and mitomycin based HDCT followed by ABMT in a preoperative setting. In our study, we evaluate clinical, histological, and molecular markers and their associations with response to treatment and survival in this highly selected patient cohort.

Patients who underwent resection survived significantly longer than unresected patients. These data demonstrate that

resectability after preoperative chemotherapy is prognostically beneficial for patients with advanced gastric cancer. However, the benefit of post-chemotherapeutic surgical treatment on the overall survival of the entire patient cohort could not be shown unequivocally, because only the patients with a clinical response underwent gastrectomy in our study.

Interestingly, the histological examination of gastrectomy specimens and biopsies obtained after chemotherapy could predict overall survival of the patients. Histological responders underwent surgical treatment in all cases and survived significantly longer than non-responders. This is in accordance with previous studies, which reported a trend towards decreased survival in patients with a poor histological response.^{44, 45} However, it has to be considered that a detailed histopathological evaluation of histological regression of gastric cancer using several paraffin wax blocks of gastrectomy specimens is not entirely comparable to the findings investigating biopsies of patients with non-resectable cancer. To minimise the bias in this comparison, five biopsies with at least 18 sections for each biopsy were analysed.

The molecular markers analysed in this study included MSI, LOH at the p53 and APC loci, expression and mutation of the p53 tumour suppressor gene, and expression of the proapoptotic protein BAX.

Overexpression of p53 and the presence of p53 mutations in exons 5–9 in pretreatment biopsy specimens were significantly associated with increased overall survival. In addition, p53 protein expression and mutation status were the only clinical or molecular parameters associated with objective tumour regression and histological response.

Given the important role of wild-type p53 in apoptosis and the association of p53 mutations with poor survival in many tumour types,⁴⁶ these results were surprising. In contrast to our results, p53 expression detected by immunohistochemistry in pretreatment endoscopic biopsies was associated with poor response to neoadjuvant cisplatin based chemotherapy in patients with gastric cancer in two studies.^{47, 48} In a study by Kubicka *et al*,⁴⁹ positive lymph nodes and p53 mutations were the only significant adverse prognostic markers for survival after curative resection for gastric cancer. However, there are

also reports in which p53 overexpression and/or mutation is associated with better outcome after treatment of bladder cancer or glioblastoma.^{50, 51}

In our study, seven of the responding tumours, but only one of the non-responding tumours, showed mutations in exons 5–9 of the p53 gene. In addition, p53 LOH was found in the gastric cancers of eight of the responders and three of the non-responders. Two of the responders had both p53 mutation and p53 LOH. Fifty per cent of gastric cancers contain p53 mutations, and p53 LOH has been reported in 26–83% of cases.⁵² Sano *et al* found both LOH and p53 mutations in more than 60% of gastric tumours.⁵³ In contrast, Kobayashi *et al* demonstrated cases of gastric cancer that displayed LOH and did not contain p53 mutations, and vice versa,⁵⁴ which is similar to our findings. Ikeguchi *et al* described cisplatin induced apoptosis more frequently in gastric cancer cell lines with a wild-type p53 gene.⁵⁵ In contrast, Grundei *et al* found that LOH at the p53 locus in advanced gastric cancer is associated with a good clinical response to cisplatin based neoadjuvant chemotherapy.⁵⁶

A possible explanation is that p53 overexpression does not correlate with the inactivation status of the gene,^{57, 58} or that we may have detected overexpression of the wild-type protein⁵⁹ in a subset of patients. This could indicate an accelerated response to stress factors (such as DNA damage induced by cisplatin⁶⁰), with subsequent effective tumour cell elimination by apoptosis. However, the association between p53 expression and mutational status on the one hand and between p53 mutations and prolonged overall survival on the other hand argues against this hypothesis.

p53 is known to be an important determinant of DNA damage induced apoptosis, and loss of p53 in tumours is associated with an unfavourable prognosis in many forms of cancer.^{61–63} Wild-type p53 is thought to render tumours more sensitive to treatment by the induction of apoptosis, and p53 inactivation may lead to resistance to treatment. However, because p53 is also responsible for prolonged cell cycle arrest after chemotherapy induced genetic damage, it is expected to facilitate DNA repair in the absence of an apoptotic response. Therefore, tumours that inactivate p53 during progression should be less capable of DNA repair and more sensitive to a DNA damage induced mitotic catastrophe. Specific mutant forms of p53 also confer a gain of function phenotype, manifested by augmented cell growth and tumorigenic potential. Blandino *et al* found that mutations at p53His175 and p53His179 provided substantial resistance to etoposide, whereas the protective effect of p53His273 and p53Trp248 was much milder.⁶⁴ Mutant p53 can inhibit apoptosis after treatment of cancer cells with low drug concentrations (for example, cisplatin), but had no effect after the presence of high concentrations. These findings suggest a possible selective gain of function of mutant p53 that depends on the particular mutation and the identity and concentration of the chemotherapeutic drug. In summary, loss of p53, and possibly gain of function of the mutant p53 protein, might not lead to increased resistance of tumours to treatment, but could be a factor that contributes to sensitivity to chemotherapy.⁶⁵

Microsatellite analyses revealed that none of the patients with advanced gastric cancers showed an MSI-H genotype. This is in agreement with the better prognosis of MSI positive gastric cancer,⁶⁶ leading to an under-representation of MSI positive tumours in this highly selected patient cohort with advanced tumours.

“p53 protein expression and mutation status were the only clinical or molecular parameters associated with objective tumour regression and histological response”

Preoperative chemotherapy can increase the rate of curative resection in locally advanced gastric carcinoma. However, a

Take home messages

- Positive p53 immunostaining, positive p53 mutation status before chemotherapy, strong histological regression induced by preoperative high dose chemotherapy (HDCT), and surgical treatment were significantly associated with response to chemotherapy and prolonged overall survival
- Thus, positive p53 immunostaining and p53 mutation status in pretreatment tumour biopsies might be useful molecular predictors of response and prognosis in patients with advanced gastric cancer treated by preoperative HDCT
- Clinical or biological markers that could predict response to such treatment would be extremely useful
- Prospective follow up evaluation and investigation of additional molecular markers are warranted to define effective predictors for response to chemotherapy in patients with gastric cancer

significant response or survival advantage is only seen in less than half of patients treated.⁶⁷ The only prognostic factors that are commonly accepted in gastric cancer are clinicopathological features, such as performance status, patient age, or macroscopic tumour type.^{68, 69} The examination of molecular alterations on pretreatment biopsy samples may be of particular importance for identifying those patients who would be suitable for preoperative chemotherapy, in an attempt to achieve resectability, the main prognostic factor for survival in gastric cancer. Without appropriate predictive markers, patient selection is based on clinical staging parameters and results in the treatment of more than twice as many patients than will eventually benefit from such treatment.

The results of our study indicate that overexpression and mutation of p53 in pretreatment biopsy specimens could predict overall survival in patients with advanced gastric cancer receiving preoperative HDCT. The availability of clinical or biological markers to predict response to such treatment would be highly desirable. Prospective follow up evaluation and investigation of additional molecular markers are warranted to define effective predictors for response to chemotherapy in patients with gastric cancer.

Authors' affiliations

F Bataille, P Rümmele, W Dietmaier, D Gaag, P Wild, F Hofstädter, A Hartmann, Department of Pathology, University of Regensburg, 93042 Regensburg, Germany
F Klebl, Department of Internal Medicine I, University of Regensburg
A Reichle, Department of Hematology and Oncology, University of Regensburg

REFERENCES

- 1 Boring CC, Squires TS, Tong T, *et al*. Cancer statistics 1994. *CA Cancer J Clin* 1994;**44**:7–26.
- 2 Siewert JR, Bottcher K, Stein HJ, *et al*. Relevant prognostic factors in gastric cancer: ten-year results of the German gastric cancer study. *Ann Surg* 1998;**228**:449–61.
- 3 Wilke H, Preusser P, Fink U, *et al*. New developments in the treatment of gastric carcinoma. *Semin Oncol* 1990;**17**(suppl 2):61–70.
- 4 Ajani JA, Roth JA, Ryan MB, *et al*. Intensive preoperative chemotherapy with colony-stimulating factor for resectable adenocarcinoma of the esophagus or gastroesophageal junction. *J Clin Oncol* 1993;**11**:22–8.
- 5 Rougier Ph, Lasser Ph, Ducreux M, *et al*. Preoperative chemotherapy of locally advanced gastric cancer. *Ann Oncol* 1994;**5**(suppl 3):59–68.
- 6 Kelsen D, Karpeh M, Schwartz G, *et al*. Neoadjuvant chemotherapy of high-risk gastric cancer: a phase II trial of preoperative FAMTX and postoperative intraperitoneal fluorouracil–cisplatin plus intravenous fluorouracil. *J Clin Oncol* 1996;**14**:1818–28.
- 7 Melcher AA, Mort D, Maughan TS. Epirubicin, cisplatin and continuous infusion 5-fluorouracil (ECF) as neoadjuvant chemotherapy in gastroesophageal cancer. *Br J Cancer* 1996;**74**:1651–4.
- 8 Kelsen DP. Adjuvant and neoadjuvant therapy for gastric cancer. *Semin Oncol* 1996;**23**:379–89.
- 9 Tanigawa N, Morimoto H. Significance of surgical adjuvant chemotherapy for gastric cancer. *J Surg Oncol* 1991;**46**:203–7.
- 10 Yamaue H, Tanimura H, Noguchi K, *et al*. Chemosensitivity testing of fresh human gastric cancer with highly purified tumor cells using the MIT assay. *Br J Cancer* 1992;**66**:794–9.

- 11 **Saikawa Y**, Kubota T, Furukawa T, *et al*. Single-cell suspension assay with an MTT end point is useful for evaluating the optimal adjuvant chemotherapy for advanced gastric cancer. *Jpn J Cancer Res* 1994;**85**:762–5.
- 12 **Loewe SW**, Ruley HE, Jacks T, *et al*. p53-dependent apoptosis modulates the cytotoxicity of anticancer agents. *Cell* 1993;**74**:957–67.
- 13 **Zamble DB**, Jacks T, Lippard SJ. p53-dependent and -independent responses to cisplatin in mouse testicular teratocarcinoma cells. *Proc Natl Acad Sci U S A* 1998;**95**:6163–8.
- 14 **Strasser A**, Harris AW, Jacks T, *et al*. DNA damage can induce apoptosis in proliferating lymphoid cells via p53-independent mechanisms inhibitable by Bcl-2. *Cell* 1994;**79**:329–39.
- 15 **Zanke BW**, Boudreau K, Rubie E, *et al*. The stress-activated protein kinase pathway mediates cell death following injury induced by cis-platinum, UV irradiation or heat. *Curr Biol* 1996;**6**:606–13.
- 16 **Sanchez-Perez I**, Murguía JR, Perona R. Cisplatin induces a persistent activation of JNK that is related to cell death. *Oncogene* 1998;**16**:533–40.
- 17 **Gonzalez VM**, Fuentes MA, Alonso C, *et al*. Is cisplatin-induced cell death always produced by apoptosis? *Mol Pharmacol* 2001;**59**:657–63.
- 18 **Eliopoulos AG**, Kerr DJ, Herod J, *et al*. The control of apoptosis and drug resistance in ovarian cancer: dependence of p53 and Bcl-2. *Oncogene* 1995;**11**:1217–28.
- 19 **Strathdee G**, MacKean MJ, Illand M, *et al*. A role for methylation of the hMLH1 promoter in loss of hMLH1 expression and drug resistance in ovarian cancer. *Oncogene* 1999;**18**:2335–41.
- 20 **Mello JA**, Acharya S, Fishel R, *et al*. The mismatch-repair protein hMSH2 binds selectively to DNA adducts of the anticancer drug cisplatin. *Chem Biol* 1996;**3**:579–89.
- 21 **Duckett DR**, Drummond JT, Murchie AI, *et al*. Human MutS- α recognized damaged DNA base pairs containing O₆-methylguanine, O₆-methylthymine or the cisplatin-d(GpG) adduct. *Proc Natl Acad Sci U S A* 1996;**93**:6443–7.
- 22 **Yamada M**, O'Regan E, Brown R, *et al*. Selective recognition of a cisplatin-DNA adduct by human mismatch repair proteins. *Nucleic Acids Res* 1997;**25**:491–6.
- 23 **Drummond JT**, Anthony A, Brown R, *et al*. Cisplatin and adriamycin resistance are associated with MutL α and mismatch repair deficiency in an ovarian tumor cell line. *J Biol Chem* 1996;**271**:19645–8.
- 24 **Aebi S**, Kurdi-Haidar B, Gordon R, *et al*. Loss of DNA mismatch repair in acquired resistance to cisplatin. *Cancer Res* 1996;**56**:3087–90.
- 25 **Vaisman A**, Varchenko M, Umar A, *et al*. The role of hMLH1, hMSH3 and hMSH6 defects in cisplatin and oxaliplatin resistance: correlation with replicative bypass of platinum-DNA adducts. *Cancer Res* 1998;**58**:3579–85.
- 26 **Fink D**, Nebel S, Norris PS, *et al*. Enrichment for DNA mismatch repair-deficient cells during treatment with cisplatin. *Int J Cancer* 1998;**77**:741–6.
- 27 **Hermanek P**. Tumors of the gastrointestinal tract and the pancreas: histopathology, staging and prognosis. *Anticancer Res* 1999;**19**:2393–6.
- 28 **Oken MM**, Creech RH, Tormey DC, *et al*. Toxicity and response criteria of the Eastern Cooperative Oncology Group. *Am J Clin Oncol* 1982;**5**:649–55.
- 29 **Lauren P**. The two histological main types of gastric carcinoma: diffuse and so-called intestinal-type carcinoma. *Acta Pathol Microbiol Scand* 1965;**64**:31–49.
- 30 **Japanese Research Society for Gastric Cancer**. *Japanese classification of gastric carcinoma*, 1st English ed. Tokyo: Kanehara, 1995.
- 31 **Jass JR**, Sobin LH, Watanabe H. The World Health Organization's histological classification of gastrointestinal tumors. A commentary on the second edition. *Cancer* 1990;**15**:66:2162–7.
- 32 **Wild P**, Knuechel R, Dietmaier W, *et al*. Laser microdissection and microsatellite analyses of breast cancer reveal a high degree of tumor heterogeneity. *Pathobiology* 2000;**68**:180–90.
- 33 **Zhang L**, Cui X, Schmitt K, *et al*. Whole genome amplification from a single cell: implications for genetic analysis. *Proc Natl Acad Sci U S A* 1992;**89**:5847–51.
- 34 **Dietmaier W**, Hartmann A, Wallinger S, *et al*. Multiple mutation analyses in single tumor cells with improved whole genome amplification. *Am J Pathol* 1999;**154**:83–95.
- 35 **Boland CR**, Thibodeau SN, Hamilton SR, *et al*. A National Cancer Institute workshop on microsatellite instability for cancer detection and familial predisposition: development of international criteria for the determination of microsatellite instability in colorectal cancer. *Cancer Res* 1998;**58**:5248–57.
- 36 **Hartmann A**, Zanardo L, Bocker-Edmonston T, *et al*. Frequent microsatellite instability in sporadic tumors of the upper urinary tract. *Cancer Res* 2002;**62**:6796–802.
- 37 **Schlegel J**, Bocker T, Zirngibl H, *et al*. Detection of microsatellite instability in human colorectal carcinomas using a non-radioactive PCR-based screening technique. *Virchows Arch* 1995;**426**:223–7.
- 38 **Hartmann A**, Rosner U, Schlake G, *et al*. Clonality and genetic divergence in multifocal low-grade superficial urothelial carcinomas as determined by chromosomes 9 and p53 deletion analysis. *Lab Invest* 2000;**80**:709–18.
- 39 **Ahlgren JD**, Macdonald JS. Gastric cancer: epidemiology, pathology, detection, and staging. In: Macdonald JS, Hill MC, Roberts IM, eds. *Gastrointestinal oncology*. Philadelphia: JB Lippincott, 1992:151–8.
- 40 **Boring CC**, Squires TS, Tong T. Cancer statistics, 1993. *C A Cancer J Clin* 1993;**43**:7–26.
- 41 **Moriguchi S**, Maehara Y, Korenaga D, *et al*. Prediction of survival time after curative surgery for advanced gastric cancer. *Eur J Surg Oncol* 1992;**18**:287–92.
- 42 **Roder JD**, Bottcher K, Siewert JR, *et al*. Prognostic factors in gastric carcinoma. *Cancer* 1993;**72**:2089–97.
- 43 **Rougier Ph**, Lasser Ph, Ducreux M, *et al*. Preoperative chemotherapy of locally advanced gastric cancer. *Ann Oncol* 1994;**5**(suppl 3):59–68.
- 44 **Yonemura Y**, Ooyama S, Matsumoto H, *et al*. Pancreaticoduodenectomy in combination with right hemicolectomy for surgical treatment of advanced gastric cancer located in the lower half of stomach. *Int Surg* 1991;**76**:226–9.
- 45 **Yonemura Y**, Katamyama K, Kamata T, *et al*. Surgical treatment of advanced gastric cancer with metastasis in para-aortic lymph node. *Int Surg* 1991;**76**:222–3.
- 46 **Erenpreisa J**, Cragg MS. Mitotic death: a mechanism of survival? A review. *Cancer Cell Int* 2002;**1**:1.
- 47 **Cascinu S**, Graziano F, Del Ferro E, *et al*. Expression of p53 protein and resistance to preoperative chemotherapy in gastric cancer. *Cancer* 1998;**83**:1917–22.
- 48 **Nakata B**, Chung KH, Ogawa M, *et al*. p53 expression as a predictor of the response to chemotherapy in gastric cancer. *Surg Today* 1998;**28**:595–8.
- 49 **Kubicka S**, Claas C, Staab S, *et al*. p53 mutation pattern and expression of c-erbB2 and c-met in gastric cancer: relation to histological subtypes, Helicobacter pylori infection, and prognosis. *Dig Dis Sci* 2002;**47**:114–21.
- 50 **Cote RJ**, Esrig D, Groshen S, *et al*. p53 and treatment of bladder cancer. *Nature* 1997;**358**:123–5.
- 51 **Tada M**, Matsumoto R, Iggo RD, *et al*. Selective sensitivity to radiation of cerebral glioblastomas harboring p53 mutations. *Cancer Res* 1998;**58**:1793–7.
- 52 **Fenoglio-Preiser CM**, Wang J, Stemmermann GN, *et al*. TP53 and gastric carcinoma: a review. *Hum Mutat* 2003;**21**:258–270.
- 53 **Sano T**, Tsujino T, Yoshida K, *et al*. Frequent loss of heterozygosity on chromosomes 1q, 5q, and 17p in human gastric carcinomas. *Cancer Res* 1991;**51**:2926–31.
- 54 **Kobayashi M**, Kawashima A, Mai M, *et al*. Analysis of chromosome 17p13 (p53 locus) alterations in gastric carcinoma cells by dual color fluorescence in situ hybridization. *Am J Pathol* 1996;**149**:1575–84.
- 55 **Ikeguchi M**, Tatebe S, Kaibara N, *et al*. Changes in levels of expression of p53 and the product of the bcl-2 in lines of gastric cancer cells during cisplatin-induced apoptosis. *Eur Surg Res* 1997;**29**:396–402.
- 56 **Grundeit T**, Vogelsang H, Ott K, *et al*. Loss of heterozygosity and microsatellite instability as predictive markers for neoadjuvant treatment in gastric carcinoma. *Clin Cancer Res* 2000;**6**:4782–8.
- 57 **Hashimoto T**, Tokuchi Y, Hayashi M, *et al*. P53 null mutations undetected by immunohistochemical staining predict a poor outcome with early-stage non-small cell lung carcinomas. *Cancer Res* 1999;**59**:5572–7.
- 58 **Hartmann A**, Blaszyk H, McGovern RM, *et al*. P53 mutations inside and outside of exons 5–8: the patterns differ in breast and other cancers. *Oncogene* 1995;**16**:681–8.
- 59 **Hartwell LH**, Kastan MB. Cell cycle control and cancer. *Science* 1994;**266**:1821–8.
- 60 **Pinto AL**, Lippard SJ. Binding of the antitumor drug cis-diaminedichloroplatinum(II) (cisplatin) to DNA. *Biochim Biophys Acta* 1985;**780**:167–80.
- 61 **Cordon-Cardo C**, Dalbagni G, Sarkis AS, *et al*. Genetic alterations associated with bladder cancer. *Important Adv Oncol* 1994;**71**–83.
- 62 **Falette N**, Paperin MP, Treilleux I, *et al*. Prognostic value of p53 gene mutations in a large series of node-negative breast cancer patients. *Cancer Res* 1998;**58**:1451–5.
- 63 **Molina R**, Segui MA, Climent MA, *et al*. p53 oncoprotein as a prognostic indicator in patients with breast cancer. *Anticancer Res* 1998;**18**:507–11.
- 64 **Blandino G**, Levine AJ, Oren M. Mutant p53 gain of function: differential effects of different p53 mutants on resistance of cultured cells to chemotherapy. *Oncogene* 1999;**18**:477–85.
- 65 **Gudkov AV**, Komarova EA. The role of p53 in determining sensitivity to radiotherapy. *Nat Rev Cancer* 2003;**3**:117–29.
- 66 **Chiaravalli AM**, Cornaggia M, Furlan D *et al*. The role of histological investigation in prognostic evaluation of advanced gastric cancer. Analysis of histological structure and molecular changes compared with invasive pattern and stage. *Virchows Arch* 2001;**439**:158–69.
- 67 **Ajani JA**, Mansfield PF, Lynch PM, *et al*. Enhanced staging and all chemotherapy preoperatively in patients with potentially resectable gastric carcinoma. *J Clin Oncol* 1999;**17**:2403–11.
- 68 **Rougier P**, Ducreux M, Mahjoubi M, *et al*. Efficacy of combined 5-fluorouracil and cisplatin in advanced gastric carcinomas: a phase II trial with prognostic factor analysis. *Eur J Cancer* 1994;**30**:1263–9.
- 69 **Cascinu S**, Labianca R, Alessandrini P, *et al*. Intensive weekly chemotherapy for advanced gastric cancer using fluorouracil, cisplatin, epi-doxorubicin, 6S-leucovorin, glutathione, and filgrastim: a report from the Italian Group for Digestive Tract Cancer. *J Clin Oncol* 1997;**15**:3313–19.